

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number  
WO 02/04617 A2

(51) International Patent Classification: C12N 9/50

(21) International Application Number: PCT/EP01/07255

(22) International Filing Date: 26 June 2001 (26.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
00114861.8 11 July 2000 (11.07.2000) EP(71) Applicant (for all designated States except US): SOCI-  
ETE DES PRODUITS NESTLE S.A. [CH/CH]; P.O. Box  
353, CH-1800 Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BUCHELI, Peter  
[CH/FR]; 73, avenue Georges Sand, F-37700 La Ville  
aux Dames (FR). LAIOL, Maryse [FR/FR]; 35-39, rue  
Jacques Cartier, F-37000 Tours (FR). MC CARTHY,  
James [US/FR]; 298 Vallée de Vantuchot, F-37210  
Noizay (FR). KOCHHAR, Sunil [IN/CH]; Chemin du  
Grammont 2, CH-1073 Savigny (CH).(74) Agent: STRAUS, Alexander; Becker, Kurig, Straus,  
Bavariastrasse 7, 80336 München (DE).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,  
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,  
ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

## Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL CACAO ENDOPROTEINASES AND THEIR USE IN THE PRODUCTION OF COCOA FLAVOUR

(57) Abstract: The present invention pertains to novel aspartic endoproteinases from Th. cacao which are involved in the production of cocoa flavour and DNA sequences coding for them. In particular, the present invention relates to the use of said enzymes in the manufacture of cocoa flavour.

WO 02/04617 A2

**Novel cacao endoproteinases and their use  
in the production of cocoa flavour**

5

The present invention pertains to novel endoproteinases involved in the production of cocoa flavour and the DNA coding for them. In particular, the present invention relates to the use of said enzymes for the manufacture of cocoa flavour.

- 10 It is known that in processing cacao beans the generation of the typical cocoa flavour requires two steps - the fermentation step, which includes air-drying of the fermented material and the roasting step. Though roasting seems to be the key stage of obtaining cocoa flavour subjecting non fermented beans to a roasting step does not yield cocoa flavour suggesting that during the fermentation step precursors are produced that are essential for flavour
- 15 generation (Rohan J. Food Sci. 29 (1964), 456-459).

- During fermentation two major activities may be observed. First, the pulp surrounding the beans is degraded by micro-organisms with the sugars contained in the pulp being largely transformed to acids, especially acetic acid (Quesnel et al. J. Sci. Food. Agric. 16 (1965),
- 20 441-447; Ostovar and Keeney, J. Food. Sci. 39 (1973), 611-617). The acids then slowly diffuse into the beans and eventually cause an acidification of the cellular material. Second, fermentation also results in a release of peptides exhibiting differing sizes and a generation of a high level of hydrophobic free amino acids. This latter finding led to the hypothesis that proteolysis occurring during the fermentation step is not due to a random protein hydrolysis
- 25 but seems to be rather based on the activity of specific endoproteinase (Kirchhoff et al., Food Chem 31 (1989), 295-311). This specific mixture of peptides and hydrophobic amino acids is deemed to represent cocoa-specific flavour precursors.

So far in cacao beans several proteolytic enzyme activities have been investigated and

checked for their putative role in the formation of cocoa flavour precursors.

An aspartic endoproteinase activity which is optimal at very low pH (pH 3.5) and is inhibited by pepstatin A has been identified. A polypeptide described to have this activity has been  
5 isolated and is described to consist of two peptides (29 and 13 kDa) which are deemed to be derived by self-digestion from a 42 kDa pro-peptide (Voigt et al., J. Plant Physiol. 145 (1995), 299-307). The enzyme cleaves protein substrates between hydrophobic amino acid residues to produce oligopeptides with hydrophobic amino acid residues at the ends (Voigt et al., Food Chem. 49 (1994), 173-180). The enzyme accumulates with the vicilin-class (7S)  
10 globulin during bean ripening. Throughout germination, its activity remains constant during the first days and does not decrease before the onset of globulin degradation (Voigt et al., J. Plant Physiol. 145 (1995), 299-307).

A cysteine endoproteinase activity had been isolated which is optimal at a pH of 5. This  
15 enzymatic activity is believed not to split native storage proteins in ungerminated seeds. Cysteine endoproteinase activity increases during the germination process when degradation of globular storage protein occurs (Biehl et al., Cocoa Research Conference, Salvador, Bahia, Brasil, 17-23 Nov. 1996).

Moreover, a carboxypeptidase activity has been identified which is inhibited by PMSF and thus belongs to the class of serine proteases. It is stable over a broad pH range with a maximum activity at pH 5.8. This enzyme does not degrade native proteins but preferentially splits hydrophobic amino acids from the carboxy-terminus of peptides. Yet, peptides with  
20 carboxy-terminal arginine, lysine, or proline residues are seemingly resistant to degradation. The rate of hydrolysis has been found to be not only determined by the carboxy-terminal  
25 amino acid as such, but also to be affected by the neighbouring amino acid residue (Bytof et al., Food Chem. 54 (1995), 15-21).

During the second step of cocoa flavour production - the roasting step - the oligopeptides and  
30 amino acids generated at the stage of fermentation have been found to obviously undergo a Maillard reaction with reducing sugars present eventually producing the substances

responsible for the cocoa flavour as such. This hypothesis has been confirmed in an experiment, wherein an oligopeptide fraction isolated after fermentation of cacao beans had been subjected to roasting in the presence of free amino acids and reducing sugars to obtain cocoa flavour (Mohr et al., Fette, Seifen, Anstrichmittel 73 (1971), 515-521 and 78 (1976), 88-95).

5

Cocoa-specific aroma has also been obtained in an experiment wherein acetone dry powder (AcDP) prepared from unfermented ripe cacao beans was subjected to autolysis at a pH of 5.2 followed by roasting in the presence of reducing sugars. It was conceived that under these conditions preferentially free hydrophobic amino acids and hydrophilic peptides should be generated and the peptide pattern thus obtained was similar to that of extracts from fermented cacao beans. An analysis of free amino acids revealed that Leu, Ala, Phe and Val were the predominant amino acids liberated in fermented beans or autolysis (Voigt et al., Food Chem. 49 (1994), 173-180). In contrast to these findings no cocoa-specific flavour could be detected when AcDP was subjected to autolysis at a pH of as low as 3.5, the pH, at which the known aspartic endoproteinase shows activity. Only few free amino acids were found to be released but a large number of hydrophobic peptides were formed. This may be explained by the aspartic endoproteinase having a high activity at this pH with the carboxypeptidase being substantially inactive under these conditions. When incubating peptides obtained after autolysis of AcDP at a pH of 3.5 with carboxypeptidase A from porcine pancreas at pH 7.5 hydrophobic amino acids were preferentially released. The pattern of free amino acids and peptides was rather similar to that found in fermented cacao beans and in the proteolysis product obtained by autolysis of AcDP at pH 5.2. After roasting of the amino acids and peptides mixture as above, a cocoa aroma could be generated. On the contrary, with a synthetic mixture of free amino acids alone whose composition was similar to the spectrum found in fermented beans cocoa flavour could not be detected after roasting, indicating that both the peptides and the amino acids are important for this purpose (Voigt et al., Food Chem. 49 (1994), 173-180).

Apart from the enzymes also the protein source of the peptides/amino acids seems to be of importance for the generation of cocoa flavour.

- During cacao bean fermentation, the percentage reduction of protein concentration observed for vicilin and albumin was 88.8% and 47.4%, respectively (Amin et al., J. Sci. Food Agric. 76 (1998), 123-128). When peptides obtained by proteolysis of the globulin fraction were post-treated with carboxypeptidase, hydrophobic amino acids (Leu, Phe, Ala, Val, Tyr) were preferentially released and a typical cocoa aroma was detected after roasting in the presence of reducing sugars (Voigt et al., Food Chem. 50 (1994), 177-184). In contrary to that, the predominant amino acids released from the albumin-derived peptides were aspartic acid, glutamic acid and asparagine. Furthermore, no cocoa aroma was detected with the albumin fraction. It was therefore concluded that cocoa-specific aroma precursors are preferentially derived from the vicilin-like globulin of cacao bean. Consequently, the mixture of hydrophobic free amino acids and remaining oligopeptides required for the generation of the typical cocoa flavour components seems to be determined by the particular chemical structure of the cacao vicilin-class globulins.
- These globulins isolated from cacao beans were also found to be efficiently degraded by pepsin (an aspartic endoproteinase) and chymotrypsin (a serine endoproteinase). Products derived from cacao globulins by successive proteolytic digestion with pepsin and carboxypeptidase A revealed a typical, but less pronounced cocoa aroma upon roasting. No cocoa aroma precursors were generated by degradation of globulins with chymotrypsin and carboxypeptidase A (Voigt et al., Food chem, 51 (1994), 7-14). Therefore, the specific mixture of oligopeptides and hydrophobic free amino acids required for the formation of the typical cocoa aroma is not only determined by the structure of the protein substrate but also dependent on the specificity of the cacao enzyme cleaving the protein.
- In view of the above data a hypothetical model for the generation of the said mixture of peptides and amino acids, i.e. the cocoa flavour precursors, during fermentation had been devised (Fig.1), wherein in a first step peptides having a hydrophobic amino acid at their end, are formed from storage proteins, which peptides are subsequently further degraded. For splitting off hydrophobic amino acids from peptides formed in a preceding step the above carboxypeptidase activity seems to be involved. Yet, for the stage of producing the said peptides having C-terminal hydrophobic amino acids, the only known enzymatic activity

which might be considered in this respect is an aspartic endoproteinase activity related to that mentioned above. It is also possible that the activity mentioned above is the result of different enzyme activities which are still unknown.

- 5    Though some aspects of cocoa flavour production have been elucidated there is still a need in the art to fully understand the processes going on, so that the manufacture of cocoa flavour may eventually be optimized.

10    An object of the present invention therefore resides in providing means to improve the formation of cocoa flavour during processing and manufacturing.

15    The above object has been solved by providing two novel aspartic endoproteinases derived from *Th. cacao* as identified by SEQ ID No 1 and SEQ ID No 2 or variants thereof obtained by substituting, deleting or adding one or more amino acids such, that the enzymatic activity thereof is essentially retained. The aspartic endoproteinases described here (termed TcAP1 and TcAP2 in the following) shall be capable to cleave the vicilin-class globulins isolated from cacao beans so that a successive degradation of the peptides by means of carboxypeptidase will result in a mixture of peptides and amino acids that yields a cocoa flavour upon a reaction with reducing sugars, i.e. upon roasting.

20    According to another embodiment the present invention provides DNA sequences coding for the respective endoproteinases. The DNA sequences may be derived according to the genetic code from the amino acid sequences as identified under SEQ ID Nos. 1 and 2 considering the wobble hypothesis, optionally taking account codon preferences of specific hosts, in which  
25    the DNA sequences shall be expressed. The skilled person may well devise appropriate DNA sequences based on the polypeptide sequences given and his own technical knowledge and understanding. According to a preferred embodiment the DNA sequences are as identified under SEQ ID No 3 (TcAP1) and SEQ ID No 4 (TcAP2), which DNA sequence may be varied by replacing, deleting or adding one or more nucleotides such, that the  
30    endoproteinases essentially retain their enzymatic activity.

The DNA sequences may be used for recombinantly preparing the aspartic endoproteinases of the present invention. To this end the DNA sequences are incorporated into a suitable expression vector, such as a plasmid or a viral vector, which comprise the common sequences, such as a promotor, a polylinker for alleviating the cloning of the DNA sequences therein, leader sequences, to direct the polypeptide produced out of the cell. The vectors will be selected based on the requirements of the system used, e.g. for an expression in *E. coli* the vectors pGEMEX, pUC-derivates, pGEX-2T, pET-derivates, pQE8 may be envisaged, which are widespread in use and are commercially available. As an example aspartic endoproteinases could be expressed into medium or on surface of lactic acid bacteria used in lactic products such as milk or yogurt.

For expressing the endoproteinases in e.g. yeast the vectors pNFF296, pY100, pPIC9K, pPICz and Ycpad1 may be utilized and for expression in animal cells the vectors pKCR, pEFBOS, cDM8 und pCEV4 as well as pSS-derivates (Kay R. et al., Science 236 (1987), 1299-1302) may be used. Moreover, for expressing the endoproteinases in plant cells, especially in cacao, the vector pAL76 or pBin19-derivates may be used and for insect cells e.g. the vector pAcSGNT-A.

The aspartic endoproteinases may be expressed in a prokaryotic or eukaryotic cell as mentioned above. It will be appreciated that the skilled person will be able to select, based on the need and his own technical skill, an appropriate expression system to achieve the desired goal. In case the endoproteinase shall simply be added to a protein mixture, such as isolated cacao vicilin-class globulins, the recombinant enzyme may be produced in a bacterial system such as *E. coli* or in yeast and applied on the protein material.

Yet, in view of the implication to increase the enzymatic activity in cacao itself a transgenic plant cell may be envisaged, wherein one or more copies of the endoproteinases, optionally coupled with a suitable and controllable promotor, have been incorporated into the genome of the plant cell. The introduction of the DNA sequence(s) may be achieved by e.g. homologous recombination of DNA stretches harboring one or more copies of the DNA sequences coding for the endoproteinases of the present invention into embryogenic calli

prepared beforehand. Since plant cells are totipotent a new transgenic cacao tree may be produced in this way the beans of which will exhibit more rapid degradation of the vicilin-class globulins when subjected to conditions of fermentation.

- 5 In consequence a transgenic plant, harboring one or more additional copy(ies) of a DNA sequence coding for the endoproteinases of the present invention is well within the scope of the present invention.

- The present endoproteinases may also be used for the manufacture of cocoa flavour by treating a suitable starting material (cacao bean, liquor or crumb), preferably vicilin-class globulins, with said endoproteinases of the present invention and concurrently or afterwards treating the material with carboxypeptidase to obtain a mixture of peptides and amino acids appropriate to act as cocoa flavour precursors. This mixture may then be subjected to "a roasting step", i.e. may be subjected to a reaction with reducing sugars to eventually obtain cocoa flavour.

- Since some of the enzymes involved in the generation of cocoa flavour are now at hand cocoa flavour may be produced artificially without having to rely on the common process of fermenting and roasting cacao beans. The present invention therefore also provides a method for generating cocoa flavor which comprises the step of subjecting a material suitable to yield cocoa flavour precursors, such as the known vicilin-class globulins, to an enzymatic degradation involving the use of the aspartic endoproteinases of the present invention.

- In particular, the present aspartic endoproteinases may be overexpressed in protein bodies of plant cells, especially seed cells, and then hydrolysis of the cellular protein material may be effected by treating such plant cells with an acidic solution.

- The present endoproteinases may also be used for hydrolyzing proteins by contacting a material of choice, such as the protein in isolated form or material containing the protein, such as e.g. food material, with an endoproteinase of the present invention and effecting hydrolysis to a desired degree. Examples for materials are dairy substances (whey protein,



and casein), wheat gluten, corn gluten, meat, egg protein and other protein containing vegetable substances not mentioned above such as proteins from oil seeds, including soybean protein and defatted soy protein.

5 In the figures,

Fig. 1 shows the theoretical production process of cocoa specific flavour precursors;

Fig. 2 shows a schematic representation of plant aspartic prepropeptides;

10

Fig. 3 schematically shows the cloning strategy for the isolation of the endoproteinase *TcAP1* cDNA;

15

Fig. 4 schematically shows the cloning strategy for the isolation of the endoproteinase *TcAP2* cDNA;

Fig. 5 shows a comparison between the different polypeptides obtained;

Fig. 6 shows a hydrophilicity Kyte Doolittle plot for both endoproteinas obtained.

20

Fig 7 shows the expression of *TcAP1a* and *TcAP2* in cacao beans of three different cacao clones;

25

Fig. 8 shows the results of a Northern blot analysis of *TcAP1a* and *TcAP2* expression in cacao beans from clone CCN51 at different maturation stages;

Fig. 9 shows the results of a Northern blot analysis of *TcAP1a* and *TcAP2* expression in cacao bean produced by cacao clone CCN51 at different germination stages;

30

Fig. 10 shows the results of a hydrolysis experiment of bovine haemoglobin by recombinant *TcAP2* protein in yeast culture medium and comparison with control strain pNFF296;

Fig. 11 shows the results of experiments determining the pH dependence of haemoglobin hydrolysis by recombinant TcAP2;

- 5 Fig. 12 shows the effect of different inhibitors on the hydrolysis of bovine haemoglobin by recombinant TcAP2;

Fig. 13 shows the analysis of most active pool (fractions 57-64) from the Sephacryl S-200 HiPrep 16/60 size exclusion column on a 10-20% Gradient SDS-PAGE Gel (Coomassie stained). In lanes 1-3, 12, 24, and 40.8 µg protein was loaded in each lane respectively. Complex denotes a putative covalent complex between AP and trypsin inhibitor fragments; TcAP2 denotes the 30.5 kDa polypeptide; 27.9 denotes the 27.9 kDa putative endochitinase; TI, trypsin inhibitor. The molecular weights of the markers are noted on the right;

- 15 Fig. 14 shows SDS-PAGE gel analysis of the reaction products after a Q Sepharose Fast Flow purified aspartic endoproteinase preparation was incubated in acid conditions for 1 minute and 7 hours. AP denotes the 30.5 kDa polypeptide; 27.9 denotes the 27.9 kDa putative endochitinase; TI, trypsin inhibitor. M, molecular weight markers (Precision, Biorad);

20

Fig. 15 shows denaturing size exclusion chromatography of the reaction products after a Q Sepharose Fast Flow purified aspartic endoproteinase was incubated in acid conditions for 1 minute and 7 hours respectively. The molecular weight size markers are: 1, ribonuclease A 13.7 kDa; 2, aprotinin 6.5 kDa; 3, substance P 1,347 Da; 4, N-benzoyl-gly-phe (hippuryl-phe) 326 Da; 5, phe 165 Da;

25

During the studies leading to the present invention two aspartic endoproteinases have been found which seem to participate in the enzymatic degradation of vicillin-class globulins in cocoa beans under the conditions of fermentation.

30

Aspartic endoproteinases as such are a widely distributed class of proteases in animals,

microbes, viruses and plants. All aspartic endoproteinases contain two aspartic residues at the active site and are active at acidic pH. In most of the aspartic endoproteinases, the catalytic aspartic residues are contained in a common Asp-Thr-Gly motif present in both lobes of the enzyme, with plant aspartic endoproteinases containing Asp-Ser-Gly at one of the sites.

5

Many aspartic proteinases have been detected or purified in monocots and dicots, which are either heterodimeric or monomeric. The sequences of corresponding genes predict that the active heterodimeric enzymes are derived from the processing of a single proprotein.

- 10 Though the genes and predicted proproteins for both monomeric and dimeric plant aspartic endoproteinases are quite similar they differ from mammalian and microbial counterparts by the presence of a 100 amino acids insert (a so called plant specific insert: PSI) which is absent in mammalian and microbial aspartic proteinases. This insert divides the protein in two regions: an amino-terminal and a carboxy-terminal region which show a relatively high
- 15 similarity to each other and to mammalian and microbial enzymes. The amino-terminal region contains the two active sites Asp-Thr-Gly (DTG) and Asp-Ser-Gly (DSG) (Fig 2). Although the positions of six cysteine residues are conserved, the PSI from different species are less homologous with each other than are the amino- and carboxy-terminal regions.

- 20 In view of this knowledge the conserved region has been utilized to obtain the nucleotide and amino acid sequence of aspartic endoproteinase (TcAP1) from cacao bean (clone ICS 95) as follows:

- A 1 kb internal fragment of the aspartic proteinase from cacao bean was amplified by RT-
- 25 PCR using degenerate oligonucleotides that had been chosen according to an alignment of known aspartic endoproteinase sequences and a selection of conserved regions. Based on the sequence of this fragment, primers were designed to amplify 5'- and 3'-end. Afterwards, a full-length cDNA (*TcAPIb*) was obtained by ligation of the 3' and 5' fragment using the *Bam*HI restriction site and another one (*TcAPIa*) was amplified using primers specific to
- 30 both extremities (Fig. 3).

*TcAPIa* and *TcAPIb* nucleotide sequences differ only by 6 base pairs. Some of these differences are also found in the partial 1 kb fragment. Three of the differences lead to amino acid changes in the encoded protein (Table 1). The molecular weight and the pI of the protein are not changed.

Table 1. Differences observed in the nucleotide sequences from the different cDNA fragments obtained by PCR and their impact on the protein sequence.

Position	1 kb fragment	TcAPIa	TcAPIb	Altered residue
318		T	A	L----M
431	<u>C</u>	<u>C</u>	T	No change
636	<u>G</u>	<u>G</u>	A	<u>A</u> ----T
764	<u>T</u>	<u>C</u>	<u>T</u>	No change
1189	<u>C</u>	T	<u>C</u>	V---- <u>A</u>
1376	<u>C</u>	<u>C</u>	T	No change

These differences may be explained by mistakes performed by polymerase enzymes during the PCR reactions. Another explanation could be that *TcAPIa* and *TcAPIb* are two different alleles from the same gene that we will name *TcAPI*. Furthermore, the 5'- and 3'-untranslated regions from *TcAPIa* and *TcAPIb* are identical. This argues rather for the presence of two alleles than for two different genes.

The cDNA sequences from *TcAPIa* (SEQ ID No 3) isolated from cacao bean (clone ICS95) is 1784 bp long. A putative initiation start codon was assigned by comparison with other plant aspartic proteinase sequences. It is located 63 bp from the 5' end. The open reading frame is broken by a stop codon (TAA) at position 1605, followed by a putative polyadenylation signal (TATAAA) at position 1625.

*TcAPIa* encodes a 514 amino acid protein with a predicted molecular weight of 56 kDa and a pI of 5.05. The protein shows a high similarity with plant aspartic endoproteases. Considering entire sequences, percent identity ranged between 59% observed with rice aspartic endoproteinase (Oryzasin A) and 87% with partial cotton sequence. A

hydrophobicity analysis (Fig. 6A) reveals that TcAP1a encodes a hydrophilic protein with a very hydrophobic N-terminal end, indicating the presence of a signal peptide. Two catalytic triads (DTG and DSG) are also present.

- 5 The nucleotide and amino acid sequence of aspartic endoproteinase (*TcAP2*) from cacao bean (clone CCN51) was obtained as follows:

A 1 kb internal fragment of the aspartic endoproteinase from cacao bean was amplified by RT-PCR using degenerate oligonucleotides selected as above. Based on the sequence of this  
10 fragment, primers were designed to amplify 5'- and 3'-end. Afterwards, a full-length cDNA (*TcAP2*) was amplified using primers specific to both extremities (Fig. 4).

The cDNA sequence from *TcAP2* (SEQ ID No 4) isolated from cacao bean (clone CCN51) is 1828 bp long. An initiation start codon is located 62 bp from the 5' end. The open reading  
15 frame is broken by a stop codon (TAA) at position 1606, followed by a putative polyadenylation signal (TATAAA) at position 1669.

*TcAP2* encodes a 514 amino acid protein with a predicted molecular weight of 56 kDa and a pI of 5.31. The protein shows a high similarity with plant aspartic endoproteinases.  
20 Considering entire sequences, percent identity ranged between 57% observed with rice aspartic endoproteinase (Oryzasin A) and 77% with partial cotton sequence. A hydrophobicity analysis (Fig 6B) reveals that TcAP2 encodes a hydrophilic protein with a very hydrophobic N-terminal end, indicating the presence of a signal peptide. Two catalytic triads (DTG and DSG) are also present.

25

The following examples illustrate the invention without limiting it to the same.

Cacao (*Theobroma cacao* L.) beans from ripe pods of clones ICS 95, CCN51 and EET95 were provided by Nestlé ex-R&D Center Quito (Ecuador). The beans were taken from the  
30 pods immediately after arrival at the laboratory (4-5 days after harvesting). The pulp and the seed coat were eliminated and the cotyledons were frozen in liquid nitrogen and stored at -

80°C until use.

### Example 1

#### Preparation of mRNA

5

Two beans were grounded in liquid nitrogen to a fine powder and extraction was directly performed with a lysis buffer containing 100 mM Tris HCl pH8, 1% SDS and 0.1M  $\beta$ -mercaptoethanol. RNA was extracted with one volume phenol/chloroform/isoamylalcohol (25/24/1) and centrifuged at 8000 rpm for 10 min at 4°C. The aqueous phase was washed  
10 three times with chloroform/isoamylalcohol (24/1). RNA was precipitated with 0.3M sodium acetate pH 5.2 in two volumes of ethanol. The RNA pellet obtained after centrifugation was resuspended in 100 mM Tris HCl pH 8 and a second precipitation with 2M lithium chloride was performed. The RNA pellet was washed with 70% ethanol and resuspended in DEPC treated water.

15

### Example 2

#### Cloning of aspartic proteinase cDNAs

20

A search for aspartic proteinase sequences in the GenBank database led to the identification of several plant sequences. A multiple alignment of these sequences revealed the presence of conserved regions, which have been used to design two degenerate oligonucleotides:

25

A sense primer, pAP0 (5'-GAYACNGGNAGYTCYAAYYTVTGG) has been synthesised according to the sequence Asp-Thr-Gly-Ser-Ser-Asn-Leu-Trp, which contains an active site (Asp-Thr-Gly) of the protein.

30

An antisense primer pAP4r (5'-CCATMAANACRTCNCMMARRATCC) has been synthesised according to the sequence Trp-Ile-Leu-Gly-Asp-Val-Phe, located in the C-terminal part of the protein.

Total RNA as prepared in example 1 was used to synthesize first strand cDNA with the

SMART PCR cDNA Synthesis Kit (Clontech, USA). Synthesis has been performed exactly as described in the kit instructions using 1 µg of total RNA and the Superscript<sup>TM</sup> II MMLV reverse transcriptase (Gibco BRL, USA). After synthesis, cDNA was used directly for PCR or kept at -20°C.

5

Specific cDNA amplification was performed with 2 µl first strand cDNA in 50 µl buffer containing 10mM Tris-HCl pH 8.8, 50mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.001 % (w/v) gelatin, 0.25 mM dNTP's, 30 pmoles of pAP0 and pAP4r primers and 5 units of Taq DNA polymerase (Stratagene, USA). Amplification was performed in a Bio-med thermocycler 60 (B. Braun).

10

A first denaturation step (94°C, 2 min) was followed by 30 cycles of denaturation (94°C, 1 min), primer annealing (40°C, 1.5 min) and extension (72°C, 2 min). The extension time was increased by 3 sec at each cycle. Amplification was ended by a final extension step (72°C, 10 min). The amplified fragment was cloned in pGEM<sup>®</sup>-T Easy vector and sequenced.

15

*TcAP1* and *TcAP2* full-length cDNAs were cloned using Rapid Amplification cDNA Ends PCR (RACE PCR). For *TcAP1*, the Marathon<sup>TM</sup> cDNA Amplification Kit (Clontech, USA) was used. Poly A+ RNA purified from total RNA (150 µg) with the Oligotex mRNA kit (QIAGEN, Germany) were used for the synthesis of double strand cDNA and a Marathon cDNA adaptor was ligated at both ends of the cDNA. These two steps have been performed according to the instructions of the Marathon<sup>TM</sup> cDNA Amplification Kit. For *TcAP2*, single strand cDNA has been synthesised from total RNA according to the SMART<sup>TM</sup> RACE cDNA Amplification Kit (Clontech, USA).

20

RACE PCR was performed with 5 µl Marathon adaptor-ligated double strand cDNA or 2.5 µl SMART single strand cDNA in 50 µl buffer containing 40 mM Tricine-KOH pH 9.2, 15 mM KOAc, 3.5 mM Mg(OAc)<sub>2</sub>, 3.75 µg/ml BSA, 0.005% Tween-20, 0.005% Nonidet-P40, 0.2 mM dNTP's, 0.2 µM of each primer and 1 µl Advantage 2 Polymerase mix (Clontech, USA). Amplification was performed via touchdown PCR, in a Bio-med thermocycler 60 (B. Braun).

30

A first denaturation step (94°C, 1 min) was followed by:

- 5 cycles including denaturation at 94°C for 30 sec and annealing/extension at 72°C for 7 min

- 5 cycles including denaturation at 94°C for 30 sec and annealing/extension at 70°C for 7 min

- 25 cycles including denaturation at 94°C for 20 sec and annealing/extension at 68°C for 7 min

For *TcAP1*, two specific primers were paired with the AP1 primer, specific to the Marathon cDNA Adaptor provided in the Marathon kit:

ICS5 for 5'RACE PCR reaction (5'GCAGCCACCAGCACAAAGTCCAG)

ICS3 3'RACE PCR reaction (5'CGGTTGGAAATGCTGTGCCTGTGTGG)

For *TcAP2*, two specific primers were paired with the UPM (Universal Primer Mix) primer that recognises the SMART sequence:

CCN5 for the 5'RACE PCR reaction (5'ATGTGTGCTTGCCCTTGTAGTGG)

CCN3 for the 3'RACE PCR reaction (5'CCGCAATGTAGATGAAGAAGCAGGTGG)

The amplified fragments were cloned in pGEM®-T Easy vector and sequenced. The sequence information obtained after the sequencing of RACE fragments was used to design new oligonucleotides in order to amplify the full length fragments:

25	TcAP1	TcAP1, sense primer (5'TCTGCTCAGCTTTTCTTGTCG)
		TcAP1r, reverse primer (5'GGATCACATGAAATCTTAAACAAAGTGC).
	TcAP2	TcAP2, sense primer (5'CTAATACGACTCACTATAGG)
		TcAP2r, reverse primer (5'ATCTGTGACTGTTGATAAAAAGC)

30 PCR reaction was performed exactly as for the amplification of 5'- and 3'-RACE fragments with one denaturation step (95°C, 1 min) followed by 35 cycles of denaturation (94°C, 30



sec), primer annealing (63°C, 1 min) and extension (72°C, 2 min). The extension time was increased by 3 sec at each cycle. Amplification was ended by a final extension step (72°C, 10 min). The amplified fragment TcAP1 and TcAP2 were cloned in pGEM®-T Easy or pGEM®-T vectors respectively and sequenced.

5

Furthermore, a cloning strategy was also used to obtain the full-length TcAP1 cDNA. 5'- and 3'-RACE fragments overlap for 200 base pairs. In this overlapping region an unique restriction site BamH I is present. Both fragments have been isolated using *BamH I* and *EcoR I* (present in the plasmid) and subcloned directly in pBS+ (Stratagene, USA) using the same restriction enzymes.

10

### Example 3

Sequencing and analysis of DNA sequences

15 cDNA sequencing has been performed according to standard techniques (Maniatis, A Laboratory Manual, Cold Spring Harbor, 1992). Sequence analysis and comparison were done using DNASTar programme. The sequences are shown under SEQ ID Nos 1 and 2.

### Example 4

20 Expression of *TcAP1a* and *TcAP2* in cacao plants

For the Northern blot total RNA was separated on 1.5 % agarose gel containing 6% formaldehyde in 20mM MOPS, 5mM NaOAC, 1mM EDTA pH 7. After electrophoresis, RNA was blotted onto nylon membranes (Applicgene) and hybridized with <sup>32</sup>P-labeled  
25 *TcAP1a* or *TcAP2* probe at 65°C in 250mM Na-phosphate buffer pH 7.2, 6.6% SDS, 1 mM EDTA and 1% BSA. Membranes were washed three times at 65°C for 30 min in 2XSSC, 0.1%SDS; in 1XSSC, 0.1% SDS and finally in 0.5XSSC, 0.1%SDS.

*TcAP1a* probe was amplified by PCR using TcAP1 and TcAP1r primers and *TcAP2* probe  
30 with the following primers:

TcAP2b: a sense primer (5'-CTATAGGGCAAGCAGTGGTAACAAC)

TcAP2br: an antisense primer (5'-TGACCTAAAGGCAAATCCTAGTTTC)

- PCR reaction was performed with 1 µl template cDNA in 50 µl buffer containing: 40 mM
- 5 Tricine-KOH pH 8.7, 15 mM KOAc, 3.5 mM Mg(OAc)<sub>2</sub>, 3.75µg/ml BSA, 0.005% Tween-20, 0.005% Nonidet-P40, 0.2 mM dNTP's, 0.2 µM of each primer and 1 µl 50X Advantage 2 polymerase Mix (Clontech, USA). Amplification was performed in a Bio-med thermocycler 60 (B. Braun). A first denaturation step (94°C, 1 min) was followed by 30 cycles of denaturation (94°C, 30 sec), primer annealing (63°C, 1.5 min) and extension (72°C, 2 min).
- 10 The extension time was increased by 3 sec at each cycle. Amplification was ended by a final extension step (72°C, 10 min).

Both fragments were purified with Strataprep PCR purification kit (Stratagène, USA) and labelled by the random priming procedure (*rediprime*<sup>TM</sup> II, Amersham Pharmacia Biotech).

15

- Northern blot analysis with RNA purified from mature cacao beans produced by different trees, CCN51, EET95 and ICS95 reveals that *TcAPIa* and *TcAP2* are both expressed in beans produced by the three different trees (Fig. 7A). However, *TcAP2* is much more strongly expressed than *TcAPIa* indicating that it might be the major aspartic endoproteinase
- 20 in cacao beans. RT-PCR experiments (Fig. 7B) are in agreement with these results. Confirmation of the idea that *TcAP2* is the major aspartic endoprotease activity in the bean is provided by the N-terminal sequencing of a purified native protein, which has the same sequence than *TcAP2*. Finally, the RT-PCR results presented in figure 7B also clearly show that both genes are expressed in leaves.

25

- Similar experiments performed with RNA purified from cacao beans at different stages of maturation (Fig. 8) confirm that *TcAPI* is less expressed than *TcAP2* in developing and mature beans. *TcAPI* and *TcAP2* expression increase slightly during maturation and decrease in mature beans. *TcAP2* is mainly expressed in early bean developmental stages suggesting
- 30 that the synthesis of new aspartic endoprotease falls as the bean matures.

During germination, the expression of *TcAP2* is relatively stable in contrary to that of *TcAP1*, which increases after a few days of germination with a maximum at days 4 and 7. A strong expression is also detected at 49 days after imbibition (Fig. 9).

## 5 Example 5

### cDNA expression in yeast heterologous system

The coding sequences of *TcAP1a* and *TcAP2* were overexpressed in the yeast heterologous system *Yarrowia lipolytica*.

10

*TcAP1a* and *TcAP2* were overexpressed under the control of a synthetic XPR2-derived promoter hp4d present on the *Yarrowia lipolytica* expression/secretion plasmid pNFF296. For both cDNA, in order to excrete the recombinant protein in the culture medium the signal sequence (first 24 amino acids, predicted as according to Nielsen et al., Protein Engineering

15

10 (1997), 1-6 was replaced by a lipase signal sequence present on the *Yarrowia lipolytica* expression/secretion plasmid pNFF296.

*TcAP1a* cloned in pGEM-T Easy was used as template for the amplification of the cDNA sequence coding for a mature protein without a putative signal sequence.

20

Two primers were used for the amplification of *TcAP1a*:

#### Primer C089

(5'-CCGGCCTCTTCGGCCGCCAAGCGAATATCCAATGAGAGATTGGTCAG)

25

primers at the 5' end of the predicted mature *TcAP1a* cDNA and introduces a *Sfi*I site allowing cloning in frame to a hybrid XPR2-lipase signal sequence present on the *Yarrowia lipolytica* expression/secretion plasmid pNFF296

#### Primer C090

30

(5'-CCGGCCCCACGTGGCCTTAGTGGTGGTGTGCAGCCTCGGCAAATCCAAC)

primers at the 3' end of the mature *TcAP1a* cDNA and introduces in-frame a 3xHIS sequence

just before the stop codon and the SfiI cloning site in front of the lipase terminator of pNFF296.

TcAP2 cDNA cloned in pGEM-T was used as template for the amplification of the sequence  
5 coding for the mature protein without a putative signal sequence.

Two primers were used for the amplification of TcAP2:

Primer C091

10 (5'-CCGGCCTCTTCGGCCGCCAAGCGAGTATCCAATGATGGGCTGGTTAG)  
primers at the 5' end of the predicted mature TcAP2 cDNA and introduces a SfiI site allowing  
cloning in frame to a hybrid XPR2-lipase signal sequence present on the Yarrowia lipolytica  
expression/secretion plasmid pNFF296.

15 PrimerC092

(5'-CCGGCCCCACGTGGCCTTAGTGGTGGTGTGCCGCCTCGGCGAAGCCGAC)  
primers at the 3' end of the mature TcAP2 cDNA and introduces in-frame a 3xHIS sequence  
just before the stop codon and the SfiI cloning site in front of the lipase terminator of  
pNFF296.

20

Amplification was performed with 1 µl of template cDNA (20 ng) in 10 mM KCl, 6 mM  
(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20 mM Tris-HCl, pH 8.0, 0.1% Triton X-100, 2 mM MgCl<sub>2</sub>, 0.2 mM of each  
dNTP, 10 µg ml<sup>-1</sup> BSA, 0.25 µM of each primers and 3 units of Pfu DNA polymerase  
(Stratagene, USA). PCR was performed in a Stratagene RoboCycler (Stratagene, USA). A  
25 first cycle (95°C-5 min, 50°C-1 min, 72°C-3 min) was followed by 30 cycles (95°C-1 min,  
50°C-1 min, 72°C-3 min) and a final cycle (95°C-1 min, 50°C-1 min, 72°C-10 min). The  
PCR products were purified using the Qiaquick PCR purification Kit (Qiagen INC., USA),  
digested with SfiI, and subsequently ligated into vector pNFF296 previously digested with  
SfiI. This ligation was used to transform *E. coli* BZ234 (Biozentrum, University of Basel,  
30 Switzerland). Constructs were selected on LB plates supplemented with 50 µg ml<sup>-1</sup>  
kanamycine, analyzed by mini plasmid-preparations plus restriction enzyme digestion and

finally by DNA sequence analysis. The resulting plasmids containing TcAP1a or TcAP2 were called pCY329 and pCY330, respectively.

The *Yarrowia lipolytica* host strain YLP3 was derived from strain polf (MatA ura3-302  
5 leu2-270 xpr2-322 axp-2 SUC2) by transforming said strain to leucine prototrophy with a 5.1  
kb SalI fragment carrying the *Yarrowia lipolytica* wild-type LEU2 gene (J.-M. Nicaud, pers.  
comm.) and selecting for LEU2 convertants. The *Yarrowia lipolytica* host strain was streaked  
on a YPD agar plate (1% Difco Bacto Yeast Extract, 2% Difco Bacto Peptone, 2% Glucose,  
2% Difco Bacto Agar) and grown overnight at 28°C. 4 ml of liquid YPD pH 4.0 (1% Difco  
10 Bacto Yeast Extract, 1% Difco Bacto Peptone, 1% Glucose, 50 mM Citrate buffer at pH 4.0)  
were inoculated with freshly grown cells of the YPD plate and grown in a tube on a rotary  
shaker (200 rpm, 28°C, 8-9 hrs). Of this preculture an adequate amount was used to inoculate  
20 ml YPD pH 4.0 in a 250 ml Erlenmeyer flask without baffles. This culture was shaken in a  
rotary shaker at 200 rpm at 28°C (over night) until a cell titration of  $10^8 \text{ ml}^{-1}$  has been  
15 reached. The cells were centrifuged for 5 min at 3000 g, washed with 10 ml of sterile water  
and re-centrifuged. The cellular pellet was suspended in 40 ml 0.1 M lithium acetate pH 6.0  
(adjusted with 10% acetic acid) and shaken in a 250 ml Erlenmeyer at 140 rpm at 28°C for  
60 minutes. The cells were again centrifuged for 5 min at 3000 g. The cellular pellet was  
suspended in 2 ml lithium acetate pH 6.0 and the competent cells were kept on ice until  
20 transformation.

One hundred microliters of competent cells were mixed with 5-20 µl plasmid linearized with  
NotI and 50 µg carrier DNA (herring sperm DNA sonicated to 100-600 bp, Promega, USA)  
in a 2 ml tube and incubated for 15 minutes at 28°C. 700 µl 40% PEG4000, 0.1 M lithium  
25 acetate pH 6.0 were added and the tubes heavily agitated at 240 rpm on a rotary shaker at  
28°C for 60 minutes. A volume of 1.2 ml of 0.1 M lithium acetate pH 6.0 was added and  
mixed. 250 µl were plated on selective agar plates (0.17% Difco Bacto Yeast Nitrogen Base  
w/o amino acid and ammonium sulfate, 1% glucose, 0.006% L-leucine, 0.1% sodium  
glutamate, 0.1% Difco Bacto Casamino Acids, 2% agar). The expression plasmid pNFF296  
30 carries a defective URA3 allele allowing for the selection of multiple integration of the

expression secretion cassette in the YLP3 host strain.

- Transformants (Ura<sup>r</sup>) were re-isolated on selective medium (0.17% Difco Bacto Yeast Nitrogen Base w/o amino acid and ammonium sulfate, 1% glucose, 0.006% L-leucine, 0.1% sodium glutamate, 0.1% Difco Bacto Casamino Acids, 2% agar). A series of clones was grown in shake-flasks to check for expression and secretion of aspartic proteinase into the culture medium.

- Small patches of cells were streaked on YPD agar plates and grown overnight at 28°C. The thin layers of grown cells were used to inoculate 50 ml DMI medium in 500 ml Erlenmeyers with 4 lateral baffles. DMI medium contains per liter: KH<sub>2</sub>PO<sub>4</sub>, 10 g; MgSO<sub>4</sub>·7H<sub>2</sub>O, 2.5 g; glucose, 20 g; Trace elements solution, 5.1 ml; Vitamins solution, 17 ml; urea, 3 g. Urea was dissolved in 15 ml water and sterile filtered. The initial pH of the medium was adjusted to 5.0. The cultures were shaken at 140 rpm on a rotary shaker at 28°C for three days. Aliquots of the cultures were centrifuged at maximum speed (3000 g) for 15 min. and the supernatant used for the determination of the aspartic endoproteinase activity.

- Aspartic endoproteinase activity was assayed at 42°C in a 900 µl reaction medium containing 0.2M sodium citrate buffer pH3.0, 10 mg/ml bovine haemoglobin and 150 µl yeast culture supernatant. To stop the reaction aliquots (80 µl) were added to an equal volume of TCA 8% and the precipitated protein removed by centrifugation at 13000 g. 20 µl supernatant were mixed to 250 µl O-phthalaldehyde (OPA) reagent (50 mM sodium tetraborate, 1 % SDS, 5.96 mM OPA (dissolved in 1 ml methanol) and 1.43 mM β-mercaptoethanol. Activity was then determined measuring OD at 340 nm and expressed in pmole leucine produced per mg protein. For this, we use the following linear equation ( $OD_{340nm} = 0.0156 \text{ pmoles} + 0.0088$ ), which was determined using a standard curve with L-leucine (0 to 80 pmoles). Protein concentration was determined by Bradford assay (Biorad).

- A strong activity could be detected in 12 independent clones transformed with the pCY330 construct (TcAP2). Further characterization of the TcAP2 recombinant protein was done using one clone named pCY330-33. Comparison of activity measurement with supernatant

from pCY330-33 and pNFF296 (control) clearly shows that no activity is detected in the control ( $1.44 \pm 0.52$  pmoles L-leucine/min/mg protein) and that hydrolysis of bovine haemoglobin occurs in presence of supernatant from pCY330-33 ( $25.8 \pm 1.45$  pmoles L-leucine/min/mg protein) (Fig. 10). This activity demonstrates clearly that active recombinant TcAP2 protein is produced by pCY330-33.

The recombinant TcAP2 endoproteinase detected in pCY330-33 hydrolyses bovine haemoglobin with an optimum at pH 3 (Fig. 11). Only slight activity could be detected for pH higher than 5.

The endoprotease activity detected in the medium of pCY330-33 (TcAP2) is completely inhibited by 2  $\mu$ M pepstatin, a specific inhibitor for aspartic endoproteinase. The pepstatin insensitive activity ( $1.91 \pm 1.26$  pmoles L-leucine/min/mg protein, 6.65%) is in the same range as that one measured for the control strain ( $2.26 \pm 1.26$  pmoles L-leucine/min/mg protein, 7.8%). Other inhibitors such as 1.10 phenanthroline (metallo proteases), DCI (serine proteases) and E64 (cysteine proteases) have no effect on TcAP2 activity (Fig. 12).

The data presented here clearly show that the culture medium in which yeast pCY330-33 was grown contained a protein able to hydrolyse bovine haemoglobin. Maximum activity at acidic pH and inhibition by pepstatin are two specific biochemical features for aspartic proteinases.

### **Example 6**

#### **Native protein purification**

Approximately 25 g of the frozen EET 95 cacao beans were ground to a fine powder using liquid nitrogen and extracted with cold acetone/water/5mM sodium ascorbate (80/20/5) according to a modified procedure of Hansen et al., J. Sci. Food Agric. 77 (1998), 273-281, to remove the majority of the fat and phenolic compounds. This procedure resulted in approximately 11.3 g of a fine acetone powder.

Acetone powder (5g) was extracted twice with 500 ml of buffer A (10 mM sodium phosphate pH 7.8, 2 mM EDTA, 10 mM sodium acetate) for 1 hour at 4°C. After centrifugation (7840g, 25 min, 4°C) the combined supernatants were made sequentially to 30% and 60% ammonium sulphate. All ammonium sulphate fractions were assayed for activity and the 60% ammonium sulphate precipitate was found to have the highest level of endoproteinase activity and was dialysed against buffer B (50 mM sodium phosphate pH 7.8, 1 mM EDTA).

Using an Akta Purifier (Pharmacia), 2 x 10 ml of dialysed 60% ammonium sulfate precipitate were loaded on a HiLoad 26/10 Q Sepharose Fast Flow column (Pharmacia) at 8-10°C. After loading, the column was washed with 5 column volumes of 20 mM Tris-HCl pH 8, then eluted with a linear gradient of 10 column volumes of the same buffer supplemented with 1 M NaCl. The flow rate of the column was 10 ml/min and 5 ml fractions were collected.

Fractions from the Q Sepharose Fast Flow column were assayed for aspartic endoproteinase activity and fractions showing the highest level of activity (#65-80) were pooled. The pooled fractions (75 ml) were concentrated to 2.2 ml using "Ultrafree Biomax" 4 ml filters (5 kDa Mw cut off), and loaded onto a Sephacryl S-200 HiPrep 16/60 size exclusion column (Pharmacia) equilibrated with 10mM Tris-HCl pH 8 and 500 mM NaCl at a flow rate of 0.5 ml/min. 1 ml fractions were collected and assayed for aspartic endoproteinase activity. The most active fractions were concentrated into three pools (#53-56, #57-64, #65-68) using "Ultrafree Biomax" filter. Protein concentration was determined with the micro BCA protein assay kit (Pierce, Inc) using BSA as a standard.

The most active pool (#57-64) with a specific activity of 1054 units/mg protein (1unit=100 ng leucine equivalent produced/min) has been subjected to SDS-PAGE. This gel (Fig. 13) shows that this fraction contains several polypeptides. N-terminal sequencing of the major bands revealed that only the 30.5 kDa band (DSEETDIVAL) corresponded exactly to the sequence of the cacao TcAP2 protein of the present invention. The other main polypeptides in the preparation were found to be putative protein body proteins. The 27.9 kDa polypeptide N-terminal sequence (TVISTYWGQNGFEGT) showed the strongest homology (76.9%) with a *Glycine max* acid chitinase III-A (accession AB007127). Thus, it is likely that the 27.9 kDa protein is an acid chitinase. The N-terminal sequence obtained for the 20.2 kDa polypeptide (ANSP) confirmed



that this band is the cacao trypsin inhibitor protein (accession X56509). In order to verify whether the endoproteinase was effectively composed of two subunits (29 and 13 kDa) (Voigt et al., J. Plant Physiol. 145 (1995), 299-307), several polypeptides smaller than 15.6 kDa were also sequenced. All the examined bands were found to be fragments of the 20.2 kDa cacao trypsin inhibitor protein and none corresponded to a putative 13 kDa of TcAP2. Furthermore, the fact that the 30.5 kDa polypeptide contains both catalytic triads (D<sup>108</sup>TG, D<sup>295</sup>SG) supports the idea that this polypeptide alone is proteolytically active. Therefore, TcAP2 is a novel monomeric aspartic endoproteinase.

#### Example 7

Characterisation of the native purified aspartic endoproteinase activity

**Inhibitor Sensitivity:** The inhibitor sensitivity of the native aspartic endoproteinase was determined in 300 µl reactions containing 200 mM sodium citrate, pH 3, 10 mg/ml bovine hemoglobin, and 5 µl of size exclusion purified pool #57-64 (2.4 µg protein/µl). The inhibitors were added to give a final concentration of 2 µM pepstatin, 2 mM 1,10 phenanthroline, 100 µM dichloroisocoumarin (DCI), 10 µM E-64. The enzyme activity was determined as described in example 5. The fact that only pepstatin A inhibits completely the activity (Table 2) confirms that the protease activity purified is an aspartic endoproteinase.

Table 2 Inhibitor sensitivity of the purified aspartic endoproteinase activity. Two replicates were done for each test.

Inhibitor	mM	Remaining Activity %
-	-	100
Pepstatin A	0.002	0%
1,10 Phenanthroline	2.0	86%
E-64	0.01	88%
DCI	0.1	90%

**Determination of the optimum pH:** The activity test performed at different pH values indicated that the purified enzyme had an optimal activity at pH 3.0 (data not shown).

**Example 8**

Analysis of the products formed when a partially purified aspartic endoproteinase preparation is incubated in acid conditions

5

To examine the peptides produced by the native cacao seed aspartic endoproteinase, a Q Sepharose Fast Flow partially purified preparation of TcAP2 (197 µg protein, 1.35 units of activity/µl; specific activity 821 units/mg protein) was incubated in acid conditions. 120 µl of the partially purified enzyme were mixed with 30 µl 1 M sodium citrate pH 3. Samples of 4 µl and 10 70 µl were taken out just before incubation at 42°C (t=1 min) and after seven hours. The 4 µl samples were put in SDS gel loading buffer for SDS-PAGE analysis. The reaction in the 70 µl samples was stopped by adding SDS to 1% final concentration, the samples were freeze-dried, solubilized with 100 µl 6M urea, 20 mM sodium phosphate pH 7, loaded on a Superdex Peptide HR 10/30 column (Amersham Pharmacia Biotech) and eluted with 6M urea, 20 mM sodium 15 phosphate pH 7 at ambient temperature.

The gel presented in Fig. 14 shows that after 7 hours, nearly all the proteins seen in the 1 min sample were substantially hydrolysed. Only two significant bands remain, one of which corresponds to a reduced amount of the 30.5 kDa cacao aspartic endoproteinase polypeptide 20 indicating an enhanced resistance of the aspartic endoproteinase towards autocatalytic degradation. When the products of the aspartic endoproteinase digestion were examined by high resolution size exclusion chromatography (Fig. 15), a significant proportion of small oligopeptides were detected, with a large percentage of the peptides having sizes ranging between 2 and 70 amino acids. This observation indicates that reacting the main cacao seed 25 aspartic endoproteinase (TcAP2) with proteins can generate a significant level of very small peptides, and thus that the action of this enzyme could generate a significant proportion of the cocoa flavor precursor peptides found in fermented cocoa beans.

30

## Claims

1. A recombinant aspartic endoproteinase as identified by SEQ ID No. 1 or a variant thereof obtained by substituting, deleting or adding one or more amino acids with the proviso that the enzymatic activity of the aspartic endoproteinase is essentially retained.
2. A recombinant aspartic endoproteinase as identified by SEQ ID No. 2 or a variant thereof obtained by substituting, deleting or adding one or more amino acids with the proviso that the enzymatic activity of the aspartic endoproteinase is essentially retained.
3. A DNA sequence coding for an aspartic endoproteinase according to claim 1.
4. The DNA sequence according to claim 3, which is identified by SEQ ID No. 3 or a variant thereof obtained by replacing, deleting or adding one or more nucleotides such that the polypeptide coded thereby is still essentially active.
5. A DNA sequence coding for an aspartic endoproteinase according to claim 2.
6. The DNA sequence according to claim 5, which is identified by SEQ ID No. 4 or a variant thereof obtained by replacing, deleting or adding one or more nucleotides such that the polypeptide coded thereby is still essentially active.
7. A vector comprising a DNA sequence according to any of the claims 3 to 6.
8. A cell containing a recombinant DNA sequence according to any of the claims 3 to 7.
9. The cell according to claim 8, which is a prokaryotic cell, an eukaryotic cell or a plant cell, preferably a cacao cell.

10. A transgenic plant, containing a cell according to claim 9.
11. Use of a DNA sequence according to any of the claims 3 to 6 for the manufacture of a cacao aspartic endoproteinase.
12. The use according to claim 11, wherein the aspartic endoproteinase is produced in a suitable cell.
13. The use according to claim 12, wherein the cell is a prokaryotic or eukaryotic host cell or a plant cell, preferably a cacao plant cell.
14. Use of an aspartic endoproteinase according to any of the claims 1 or 2 for the preparation of cocoa flavour.
15. Use of an aspartic endoproteinase according to any of the claims 1 or 2 for hydrolyzing proteins.
16. The use according to claim 15, wherein the proteins are derived from food material.
17. A process for producing cocoa flavour comprising subjecting a material suitable to yield cocoa flavour precursors to an enzymatic degradation, involving the use of an aspartic endoproteinase according to any of the claims 1 or 2.
18. A product containing cocoa flavour, obtainable according to the method of claim 17.
19. A process of hydrolyzing proteinaceous material in a plant comprising expressing an aspartic endoproteinase according to any of claim 1 or 2 in plant cells, especially seed cells, and then effecting hydrolysis of the cellular protein by treating such plant cells with an acidic solution.




Fig. 1: Proteolytic formation of the cocoa-specific aroma





Fig. 2: Schematic representation of plant aspartic prepropeptides.



**Fig. 3:** Cloning strategy used for the isolation of cDNA encoding aspartic endoproteinase from *Theobroma cacao*,




Fig. 4: Cloning strategy used for the isolation of cDNA encoding aspartic endoproteinase from *Theobroma cacao*,





FIG. 5



A

Hydrophilicity Plot-kyte-Doolittle for TcAP1a



B

Hydrophilicity Plot-kyte-Doolittle for TcAP2




Fig. 6: Hydrophilicity Plot-Kyte-Doolittle for the TcAP1a (A) and TcAP2 (B) sequences.





Fig. 7: *TcAP1a* and *TcAP2* expression in cacao bean produced by three different cacao clones



**Fig. 8:** Northern blot analysis of *TcAP1a* and *TcAP2*





Fig. 9: Northern blot analysis of *TcAP1a* and *TcAP2* expression in cacao bean



**Figure 10:** Hydrolysis of bovine haemoglobin by recombinant TcAP2 protein in yeast culture medium and comparison with control strain pNFF296.





Figure 11: pH dependence of haemoglobin hydrolysis by recombinant TcAP2.



**Figure 12:** Effect of different inhibitors on the hydrolysis of bovine haemoglobin by recombinant TeAP2




Fig. 13






Fig. 14




Fig. 15

## SEQUENCE LISTING

<110> Société des Produits Nestlé S.A.

<120> Novel cacao endoproteinases and their use in the  
production of cocoa flavour

<130> 80255

<140>

<141>

<150> EP 00114861.8

<151> 2000-07-11

<160> 32

<170> PatentIn Ver. 2.1

<210> 1

<211> 514

<212> PRT

<213> Theobroma cacao

<400> 1

Met Gly Arg Ile Val Lys Thr Thr Thr Val Thr Leu Phe Leu Cys Leu  
1 5 10 15

Leu Leu Phe Pro Ile Val Phe Ser Ile Ser Asn Glu Arg Leu Val Arg  
20 25 30

Ile Gly Leu Lys Lys Arg Lys Phe Asp Gln Asn Tyr Arg Leu Ala Ala  
35 40 45

His Leu Asp Ser Lys Glu Arg Glu Ala Phe Arg Ala Ser Leu Lys Lys  
50 55 60

Tyr Arg Leu Gln Gly Asn Leu Gln Glu Ser Glu Asp Ile Asp Ile Val  
65 70 75 80

Ala Leu Lys Asn Tyr Leu Asp Ala Gln Tyr Phe Gly Glu Ile Gly Ile  
85 90 95

Gly Thr Pro Pro Gln Asn Phe Thr Val Ile Phe Asp Thr Gly Ser Ser  
100 105 110

Asn Leu Trp Val Pro Ser Ser Lys Cys Tyr Phe Ser Ile Ala Cys Tyr  
115 120 125

Leu His Ser Arg Tyr Lys Ser Ser Arg Ser Ser Thr Tyr Lys Ala Asn  
 130 135 140

Gly Lys Pro Ala Asp Ile Gln Tyr Gly Thr Gly Ala Ile Ser Gly Phe  
 145 150 155 160

Phe Ser Glu Asp Asn Val Gln Val Gly Asp Leu Val Val Lys Asn Gln  
 165 170 175

Glu Phe Ile Glu Ala Thr Arg Glu Pro Ser Ile Thr Phe Leu Val Ala  
 180 185 190

Lys Phe Asp Gly Ile Leu Gly Leu Gly Phe Gln Glu Ile Ser Val Gly  
 195 200 205

Asn Ala Val Pro Val Trp Tyr Asn Met Val Asn Gln Gly Leu Val Lys  
 210 215 220

Glu Pro Val Phe Ser Phe Trp Phe Asn Arg Asp Pro Glu Asp Asp Ile  
 225 230 235 240

Gly Gly Glu Val Val Phe Gly Gly Met Asp Pro Lys His Phe Lys Gly  
 245 250 255

Asp His Thr Tyr Val Pro Ile Thr Arg Lys Gly Tyr Trp Gln Phe Asp  
 260 265 270

Met Gly Asp Val Leu Ile Gly Asn Gln Thr Thr Gly Leu Cys Ala Gly  
 275 280 285

Gly Cys Ser Ala Ile Ala Asp Ser Gly Thr Ser Leu Ile Thr Gly Pro  
 290 295 300

Thr Ala Ile Ile Ala Gln Val Asn His Ala Ile Gly Ala Ser Gly Val  
 305 310 315 320

Val Ser Gln Glu Cys Lys Thr Val Val Ser Gln Tyr Gly Glu Thr Ile  
 325 330 335

Ile Asp Met Leu Leu Ser Lys Asp Gln Pro Leu Lys Ile Cys Ser Gln  
 340 345 350

Ile Gly Leu Cys Thr Phe Asp Gly Thr Arg Gly Val Ser Thr Gly Ile  
 355 360 365

Glu Ser Val Val His Glu Asn Val Gly Lys Ala Thr Gly Asp Leu His  
 370 375 380

Asp Ala Met Cys Ser Thr Cys Glu Met Thr Val Ile Trp Met Gln Asn  
 385 390 395 400

Gln Leu Lys Gln Asn Gln Thr Gln Glu Arg Ile Leu Glu Tyr Ile Asn  
 405 410 415

Glu Leu Cys Asp Arg Leu Pro Ser Pro Met Gly Glu Ser Ala Val Asp  
 420 425 430

Cys Ser Ser Leu Ser Thr Met Pro Asn Val Ser Phe Thr Ile Gly Gly  
 435 440 445

Lys Ile Phe Glu Leu Ser Pro Glu Gln Tyr Val Leu Lys Val Gly Glu  
 450 455 460

Gly Asp Val Ala Gln Cys Leu Ser Gly Phe Thr Ala Leu Asp Val Pro  
 465 470 475 480

Pro Pro Arg Gly Pro Leu Trp Ile Leu Gly Asp Val Phe Met Gly Gln  
 485 490 495

Phe His Thr Val Phe Asp Tyr Gly Asn Leu Gln Val Gly Phe Ala Glu  
 500 505 510

Ala Ala

<210> 2

<211> 514

<212> PRT

<213> Theobroma cacao

<400> 2

Met Gly Thr Thr Ile Lys Val Val Val Leu Ser Leu Phe Ile Ser Ser  
 1 5 10 15

Leu Leu Phe Ser Val Val Ser Ser Val Ser Asn Asp Gly Leu Val Arg  
 20 25 30

Ile Gly Leu Lys Lys Met Lys Leu Asp Pro Asn Asn Arg Leu Ala Ala  
 35 40 45

Arg Leu Asp Ser Lys Asp Gly Glu Ala Leu Arg Ala Phe Ile Lys Lys  
 50 55 60

Tyr Arg Phe Arg Asn Asn Leu Gly Asp Ser Glu Glu Thr Asp Ile Val

65		70		75		80
Ala Leu Lys Asn Tyr Met Asp Ala Gln Tyr Tyr Gly Glu Ile Gly Ile						
	85		90		95	
Gly Thr Pro Thr Gln Lys Phe Thr Val Ile Phe Asp Thr Gly Ser Ser						
	100		105		110	
Asn Leu Trp Val Ser Ser Thr Lys Cys Tyr Phe Ser Val Ala Cys Tyr						
	115		120		125	
Phe His Glu Lys Tyr Lys Ala Ser Asp Ser Ser Thr Tyr Lys Lys Asp						
	130		135		140	
Gly Lys Pro Ala Ser Ile Gln Tyr Gly Thr Gly Ala Ile Ser Gly Phe						
	145		150		155	
Phe Ser Tyr Asp His Val Gln Val Gly Asp Leu Val Val Lys Asp Gln						
	165		170		175	
Glu Phe Ile Glu Ala Thr Lys Glu Pro Gly Leu Thr Phe Met Val Ala						
	180		185		190	
Lys Phe Asp Gly Ile Leu Gly Leu Gly Phe Lys Glu Ile Ser Val Gly						
	195		200		205	
Asp Ala Val Pro Val Trp Tyr Asn Met Ile Lys Gln Gly Leu Ile Lys						
	210		215		220	
Glu Pro Val Phe Ser Phe Trp Leu Asn Arg Asn Val Asp Glu Glu Ala						
	225		230		235	
Gly Gly Glu Ile Val Phe Gly Gly Val Asp Pro Asn His Tyr Lys Gly						
	245		250		255	
Lys His Thr Tyr Val Pro Val Thr Gln Lys Gly Tyr Trp Gln Phe Asp						
	260		265		270	
Met Gly Asp Val Leu Ile Ala Asp Lys Pro Thr Gly Tyr Cys Ala Gly						
	275		280		285	
Ser Cys Ala Ala Ile Ala Asp Ser Gly Thr Ser Leu Leu Ala Gly Pro						
	290		295		300	
Ser Thr Val Ile Thr Met Ile Asn His Ala Ile Gly Ala Thr Gly Val						
	305		310		315	
Val Ser Gln Glu Cys Lys Ala Val Val Gln Gln Tyr Gly Arg Thr Ile						

	325		330		335
Ile Asp Leu Leu Ile Ala Glu Ala Gln Pro Gln Lys Ile Cys Ser Gln					
340		345		350	
Ile Gly Leu Cys Thr Phe Asn Gly Ala His Gly Val Ser Thr Gly Ile					
355		360		365	
Glu Ser Val Val Asp Glu Ser Asn Gly Lys Ser Ser Gly Val Leu Arg					
370		375		380	
Asp Ala Met Cys Pro Ala Cys Glu Met Ala Val Val Trp Met Gln Asn					
385		390		395	400
Gln Val Arg Gln Asn Gln Thr Gln Asp Arg Ile Leu Ser Tyr Val Asn					
405		410		415	
Glu Leu Cys Asp Arg Val Pro Asn Pro Met Gly Glu Ser Ala Val Asp					
420		425		430	
Cys Gly Ser Leu Ser Ser Met Pro Thr Ile Ser Phe Thr Ile Gly Gly					
435		440		445	
Lys Val Phe Asp Leu Thr Pro Glu Glu Tyr Ile Leu Lys Val Gly Glu					
450		455		460	
Gly Ser Glu Ala Gln Cys Ile Ser Gly Phe Thr Ala Leu Asp Ile Pro					
465		470		475	480
Pro Pro Arg Gly Pro Leu Trp Ile Leu Gly Asp Ile Phe Met Gly Arg					
485		490		495	
Tyr His Thr Val Phe Asp Phe Gly Lys Leu Arg Val Gly Phe Ala Glu					
500		505		510	
Ala Ala					

&lt;210&gt; 3

&lt;211&gt; 1784

&lt;212&gt; DNA

&lt;213&gt; Theobroma cacao

&lt;400&gt; 3

tctgtctcgc tttttctgtc gaaatcatca ctaaaaccat ttgcgggactt gcagttatca 60  
 gaatggggag aatagtcaaa actactacag tcaactcttt tctttgtctt cttctgtttc 120  
 ctatcgtatt ttccatatcc aatgagagat tggtcagaat tggactgaaa aagagaaagt 180

```

tcgatcaaaa ctatcggttg gctgcccacc ttgattccaa ggagagagag gcatttagag 240
ctctctctaa aaagtatcgt ctccaaggga acttacaaga gtctgaggac attgatattg 300
tggcaactaaa gaactacttg gatgctcagt actttggtga gattggtatt ggcacacctc 360
cacagaactt cactgtgatt tttagacactg gtagtcttaa ttgtgtggct ccttcactta 420
agtgcatttt ctogatagct tgctatctcc attcaagata taaatcaagc cgttcaagca 480
cctacaaggc taatggtaaa ccagccgata tccaatacgg gactggagct atttctggat 540
tcttttagtg ggacaatgta caagttgggt atcttgtagt taaaaatcag gaatttatcg 600
aggcaacaag ggagcccagc ataacatttt tgggtggcaa gtttgatggg atacttggac 660
ttggatttca agagatttct gttggaaatg ctgtgootgt gtggtaacaat atgggtcaatc 720
aaggctctgt taagggaact gttttctcat ttgtggttaa ccgcgatcct gaggatgata 780
taggtgggga agttgttttt ggtggaatgg atccaaaaca ttccaagggg gatcacactt 840
acgttctcat aacgcggaaa ggatactggc agtttgatat ggggtgatgc ctgattggta 900
accaaaacac tggactttgt gctggtggct gcagtgcaat tgctgattct gggacttctc 960
tgataaccgg tctacgggct attattgctc aagtcaatca tgctattgga gcatcagggg 1020
ttgtaagtca agaatgcaag actgtagttt cacagtatgg agagacaata attgatattg 1080
ttttatctaa ggaccaacca ctgaaaattt gctcacaat aggtttgtgc acatttgatg 1140
gaactcgagg tgtaagtacg gggattgaaa gtgttgtgca tgagaatggt gggaaagcca 1200
ctggtgtatt gcattgatga atgtgttcta ctgtgagat gacagttata tggatgcaa 1260
accagcttaa gcagaaccag acacaggagc gtatacttga gtacatcaat gagctctgtg 1320
atcggttgcc tagtccaatg ggagaatcag ctggttatgg tagcagtcata tctacaatgc 1380
ctaattgtct gtccaattt ggtggaaaga tatttgagct cagccccagc cagtatgtcc 1440
tgaagatggg tgagggagat gtatgtcaat gctcagtggt attcactgct ctggatgtgc 1500
caactctctg tggacctctg tggatcttgg ggcagctctt tatggggcac ttccatacag 1560
tatttgacta tggcaacctg caagttggat ttgcccaggc tgcataagtg aaactttctg 1620
ctttataaaa caacttcatg ttatgcagtg ctatgtagac ccttagaact gtggggatta 1680
agtatcaaat gataattgca tgtaaatatc tatgcaaac tgatctgtga tcttcaactg 1740
atcggttagt gtagtgcact ttgtttaaga atttcaatgtg atcc 1784

```

&lt;210&gt; 4

&lt;211&gt; 1828

&lt;212&gt; DNA

&lt;213&gt; Theobroma cacao

&lt;400&gt; 4

```

gaccacacttt cctcttttct ttgtttgact tgcgcaaggt ggtttogaca tttcgggttaa 60
tatgggaacg actatacaag tggttgtgct gtgcctgttc atctcgtccc tcttgttttc 120
tgtgttatct tctgtatoca atgatgggct ggttagaact ggcctgaaaa agatgaaact 180
ggatccaaat aacgggctcg ctgcccggct tgactccaag gacggagagg cctcagagc 240
attcattaaa aagtatcggt tccgtaataa ctttggagac tctgaggaga ctgatatcgt 300
tgcaactaaa aactacatgg atgctcagta ctatggcgag attggtattg gaactccaac 360
acaaaagtgc actgtgatat ttgacacagg aagctcaaat ctgtgggtat catcaaccaa 420
gtgctatttc tgggttgcat gttatttoca cgagaagtac aaggcaagct attcaagtac 480
ctataagaag gatgggaaac ctgcttctat tcagtatggc actggagcta tttctggttt 540
ctttagttat gaccatgttc aagttgttga ctggtttgtg aaagatcagg aatttatgta 600
ggctactaag gagccaggct ttacatttat ggtggccaaa ttgtatggga tattaggact 660
tgggttcaag gagatttcag ttggggatgc tgtccagtg tggtaacaa tgattataca 720
aggtcttata aaggaaccag tattttcatt ttggcttaac cgaatgtgag atgaagaagc 780

```



```

aggTggtgaa attgtttttg ggggggttga tccaaaccac tacaagggca agcacacata 840
tggtcctgta actcagaaag gctactggca gtttgacatg ggtgatgttc ttattgtgta 900
caaaccaact ggatattgtg ctggcagctg tgcgcgaatt gcagattctg gaaattcttt 960
gtcggcaggt ccatcgactg tgattaccat gattaaccat gcaattggag coactggagt 1020
ggttagccag gagtgcaagg cagtggttca acaatatggg cgaaccatca ttgatttact 1080
tatagctgag gcacaacctc agaagatctg ctcccaattt ggattgtgca cttttaatgg 1140
tgctcatggt gttagcacgg goattgagag tgtggtggat gagagcaatg gaaaatcaco 1200
tggagttctt cgtgatgcta tgtgccttgc ttgtgagatg gcagtttgtt ggatgcagaa 1260
ccaagtaagg cagaatcaga ctcaagacgg catattgagc taactaaatg agctttgtga 1320
tcgggtgcga aacccaatgg gagaatctgc tgttgactgc ggaagtcttt cttccatgcc 1380
tactatttcc ttcactattg gtggcaaagt ttttgacctc actccagaag agtatattct 1440
caaggtgggt gaaggtttct aagcacagtg catcagtggt tttactgott tggatattcc 1500
tcctcctcgt ggacctctct ggattctggg agatatcttc atgggtctgt accacaccgt 1560
ctttgatttc ggtaaaactg gactcggctt cgcgcaggcg gcataaaaga tctaccaggg 1620
ggaccocagt ttttagttgt ccaccaacta ttatgtttat tgtaacttta taaagatgga 1680
ggaatcagcc taaaatctgt ctgtgtgttg cttgtaataa ttccgcocct ttgctctgtt 1740
ctagaaacta ggatttgcct ttaggtcaaa gttgtcaaaa accaagtggag aaactgtgtg 1800
ctttgctttt tatcaacagt cacagata
1828

```

&lt;210&gt; 5

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 5

gagacnggna gytcyayyt vt

22

&lt;210&gt; 6

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 6

ccatmaaac rtcnccmarx atcc

24

&lt;210&gt; 7

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 7

gcagccacca gcacaaagtg gag

23

&lt;210&gt; 8

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 8

cgggttggaa atgctgtgcc tgtgtgg

27

&lt;210&gt; 9

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 9

atgtgtgctt gccctttag tag

23

&lt;210&gt; 10

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 10

ccgcaatgta gatgaagaag caggtgg

27

&lt;210&gt; 11

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 11

tctgtctcagc ttttcttgtc g

21

&lt;210&gt; 12

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 12

ggatcacatg aaaattctta aacaaagtgc

30

&lt;210&gt; 13

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 13

ctaatacgac tcactatagg

20

&lt;210&gt; 14

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:artificial

&lt;400&gt; 14

atctgtgact gttgataaaa agc

23

&lt;210&gt; 15

&lt;211&gt; 8

&lt;212&gt; FRT

&lt;213&gt; Theobroma cacao

&lt;400&gt; 15

Asp Thr Gly Ser Ser Asn Leu Trp  
1 5

&lt;210&gt; 16

&lt;211&gt; 7

&lt;212&gt; PRT

&lt;213&gt; Theobroma cacao

&lt;400&gt; 16

Trp Ile Leu Gly Asp Val Phe  
1 5

&lt;210&gt; 17

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 17

gayacnggna gytcyayyt vtgg

24

&lt;210&gt; 18

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 18

ccatmaanac rtcnccmarr atcc

24

&lt;210&gt; 19

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

<400> 19  
gcagccacca gcacaaagtc cag

23

<210> 20  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial

<400> 20  
atgtgtggtt gccctttagt tgg

23

<210> 21  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial

<400> 21  
ccgcaatgta gatgaagaag caggtgg

27

<210> 22  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial

<400> 22  
cggttggaat tgctgtgcct gtgtgg

26

<210> 23  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: artificial

&lt;400&gt; 23

totgctcagc ttttcttctg c

21

&lt;210&gt; 24

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 24

ggatcacatg aaattcttaa acaaagtgc

29

&lt;210&gt; 25

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 25

ctaatacgac tcaactatag

20

&lt;210&gt; 26

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 26

atctgtgact gttgataaaa agc

23

&lt;210&gt; 27

&lt;211&gt; 25

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 27

ctatagggca agcagtggta acaac

25

&lt;210&gt; 28

&lt;211&gt; 25

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 28

tgacctaaaag gcaaatccta gtttc

25

&lt;210&gt; 29

&lt;211&gt; 47

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 29

ccggcctctt cggcgcccaa gcgaatatcc aatgagagat tggtcag

47

&lt;210&gt; 30

&lt;211&gt; 48

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 30

ccggccccag tggccttagt ggtggtgtgc agcctcggca aatccaac

48

&lt;210&gt; 31

&lt;211&gt; 47

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 31

ccggcctcctt cggccgcca gcgagtatoc aatgatgggc tggtttag

47

&lt;210&gt; 32

&lt;211&gt; 48

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: artificial

&lt;400&gt; 32

ccggcccaag tggccttagt ggtggtgtgc cgcctcggcg aagccgac

48



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number  
**WO 02/04617 A3**

(51) International Patent Classification: **C12N 15/57**,  
9/50, A01H 5/00, C12P 21/06, A23G 1/02

(21) International Application Number: PCT/EP01/07255

(22) International Filing Date: 26 June 2001 (26.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
00114861.8 11 July 2000 (11.07.2000) EP

(71) Applicant (for all designated States except US): **SOCI-  
ETE DES PRODUITS NESTLE S.A.** [CH/CH]; P.O. Box  
353, CH-1800 Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BUCHELI, Peter**  
[CH/FR]; 73, avenue Georges Sand, F-37700 La Ville  
aux Dames (FR). **LALOI, Maryse** [FR/FR]; 35-39, rue  
Jacques Cartier, F-37000 Tours (FR). **MC CARTHY,**  
**James** [US/FR]; 298 Vallée de Vautruchot, F-37210  
Noizay (FR).

(74) Agent: **STRAUS, Alexander**; Becker, Kurig, Straus,  
Bavariastrasse 7, 80336 München (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,  
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,  
ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(88) Date of publication of the international search report:  
13 June 2002

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

WO 02/04617 A3

(54) Title: NOVEL CACAO ENDOPROTEINASES AND THEIR USE IN THE PRODUCTION OF COCOA FLAVOUR

(57) Abstract: The present invention pertains to novel aspartic endoproteinases from *Th. cacao* which are involved in the production of cocoa flavour and DNA sequences coding for them. In particular, the present invention relates to the use of said enzymes in the manufacture of cocoa flavour.

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/EP 01/07255

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C12N15/57 C12N9/50 A01H5/00 C12P21/06 A23G1/02		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N A23G A01H C12P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
BIOSIS, WPI Data, SEQUENCE SEARCH		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	VOIGT J ET AL: "Developmental stage-dependent variation of globular storage protein and aspartic endoprotease during ripening and germination of Theobroma cacao L. seeds." JOURNAL OF PLANT PHYSIOLOGY, vol. 145, no. 3, 1994, pages 299-307, XP000972681 ISSN: 0176-1617 page 300, right-hand column, paragraph 2 --- -/--	1,2
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C <input type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
3 April 2002		17/04/2002
Name and mailing address of the ISA European Patent Office, P.B. 5616 Patentlaan 2 NL - 2260 HV Rijswijk Tel (+31-70) 340-2040, Tx: 31 651 epo nl Fax (+31-70) 340-3016		Authorized officer  Van der Schaal, C

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 01/07255

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HIRAIWA NAGAKO ET AL: "An aspartic endopeptidase is involved in the breakdown of propeptides of storage proteins in protein-storage vacuoles of plants." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 246, no. 1, 1997, pages 133-141, XP000971170 ISSN: 0014-2956 ---	
A	EMBL DATABASE Accession no U61396, Sequence identity VU061396 17 July 1996 D'ARCY-LAMETA A ET AL: "Molecular cloning and nucleotide sequence of a cDNA encoding an aspartic proteinase from Vigna unguiculata leaves" XP002155324 ---	
A	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1995 VOIGT J ET AL: "Precursors of the cocoa-specific aroma components are derived from the vicilin-class (7S) globulin of the cocoa seeds by proteolytic processing." Database accession no. PREV199698592431 XP002194847 abstract & BOTANICA ACTA, vol. 108, no. 4, 1995, pages 283-289, ISSN: 0932-8629 -----	14-19